TIC Guidelines for COVID-19

These guidelines are set up for our as well as the patients safety to prevent spread of COVID-19.

- 1. No patients with flu like symptoms or the following symptoms will be allowed in the clinic:
 - a. Fever
 - b. Cough
 - c. Sore throat
- 2. Postponing travel and face-to-face meetings deemed non-essential:
 - a. Reschedule all patients who are asymptomatic and are coming in for a follow up appointment to May if they prefer.
 - b. Reschedule all patients who have traveled out of the country in the last month.
 - c. Reschedule all pacemaker checks if no battery change is needed
 - d. Reschedule all INR Checks if previous checkup was within normal limits.
 - e. All medication refills will need to be done over the phone
- 3. Encouraging the use of Telemedicine and teleconferencing.
- 4. Have patients wear masks from home, if available, to the clinic.
- 5. Maintaining social distancing throughout the clinic:
 - a. Please keep a 6 feet distance from everybody in the clinic.
- 6. Suspending handshaking and other personal contact during daily interactions
- 7. Regular cleaning and disinfecting of workspaces and communal areas:
 - a. All staff must disinfect break room if used for lunch break
- 8. Reducing/discouraging visits from external suppliers, vendors, and other visitors
- 9. Minimizing the movement of external workers required in the clinic, such as truck drivers, delivery people, and service technicians.
- 10.Discouraging/preventing personnel from reporting to work if they are ill

- 11.Please change gloves after every patient interaction:
 - a. Scrubs should be washed and changed if in contact with patient who is suspected to have COVID-19
 - b. Please wear lab coat/gown when interacting with patients.
 - i. Change lab coat/gown if patient is suspected to have COVID-19.
- 12.Please limit exposure to patients as much as possible.

Please wash and sanitize hands at all times